

AD-A066 707 WASHINGTON UNIV SEATTLE DEPT OF CHEMISTRY F/6 7/4
COLLISIONAL RELAXATION OF NON-EQUILIBRIUM VIBRATIONAL ENERGY DI--ETC(U)
MAR 79 D F KELLEY, B D BARTON, L ZALOTAI N00014-75-C-0690
UNCLASSIFIED NR092-549-TR09 NL

| OF |
AD A066707



END
DATE
FILMED
5-79
DDC

ADA066707

LEVEL ✓

(12)

Collisional Relaxation of Non-Equilibrium Vibrational Energy
Distributions in a Thermal Unimolecular System. Surface Collisions

D. F. Kelley, B. D. Barton, L. Zalotai
and B. S. Rabinovitch

Department of Chemistry, BG-10
University of Washington
Seattle, WA 98195

Prepared for publication in J. Chem. Phys.

Technical Report No. NR092-549-TR09

Contract N00014-75-C-0690, NR092-549

DDC FILE COPY

March 15, 1979



OFFICE OF NAVAL RESEARCH
Department of the Navy
Code 473
800 N. Quincy
Arlington, VA 2217

Reproduction in whole or in part is permitted for any purpose of
the United States Government. This document has been approved for public
release; its distribution is unlimited.

79 04 02 042

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <i>(14) NR092-549-TR09</i>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Collisional Relaxation of Non-Equilibrium Vibrational Energy Distributions in a Thermal Unimolecular System. Surface Collisions.		5. TYPE OF REPORT & PERIOD COVERED <i>(17) Technical Rep's</i>
6. AUTHOR/s <i>(10) D. F. Kelley, B. D. Barton, L. Zalotai and B. S. Rabinovitch</i>		7. PERFORMING ORG. REPORT NUMBER
8. CONTRACT OR GRANT NUMBER(s) <i>(15) N00014-75-C-0690 NR092-549</i>		9. PERFORMANCE ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
10. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research, Code 743 Department of the Navy 800 N. Quincy Arlington, VA 22217		11. REPORT DATE <i>(11) 15 March 1979</i>
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES <i>(12) 9 p.</i>
		14. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES To be submitted to J. Chem. Phys.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Cyclopropane Transients Energy Transfer Unimolecular Reaction High Temperature Vibrational Energy Surfaces Variable Encounter		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Measurements are made for the first time (on cyclopropane) of the dynamics of collisional relaxation of vibrational distributions of reacting molecules at high energies, and at high temperatures. This is the same as the determination of the mean first passage time τ of molecules with respect to an absorbing barrier. A simple technique, called the Variable Encounter Method has been devised for this purpose. tau 370274B		

Collisional Relaxation of Non-Equilibrium Vibrational Energy Distributions
in a Thermal Unimolecular System.^a Surface Collisions.

by D. F. Kelley, B. D. Barton, L. Zalotai^b and B. S. Rabinovitch

Department of Chemistry, University of Washington, Seattle, WA 98195

There have been no measurements to date of the dynamics of collisional relaxation of vibrational distributions characteristic of reacting polyatomic molecules at high energies. Formal theoretical treatments have been given.^{1,2} The unknown desideratum has been the appropriate matrix of the collisional transition probabilities, P . The same matter appears in the form of the mean first passage time, $\bar{\tau}$, of molecules with respect to an absorbing barrier at energy, E_0 . Kim³ and Widom⁴ have given expressions for $\bar{\tau}$ in terms of P . For a thermal unimolecular system, the concept of an absorbing barrier corresponds to the limiting low pressure behavior. For the trivial case of a constant temperature, steady-state system, $\bar{\tau} = 1/k_0$, where k_0 is the low pressure unimolecular rate constant. But $\bar{\tau}$ has not been measured for the dynamical case of transients wherein some initial input distribution, not necessarily Boltzmann, relaxes to a final distribution characteristic of the bath temperature, whether by homogeneous or heterogeneous collision processes. Based on a quasi-accommodation model for energy transfer,⁵ as well as experimental results with bath molecules of increasing complexity,⁶ one anticipates that the wall may be at least as efficient as any bath molecule.

A very simple technique for studying the dynamics of energy transfer at a surface has been developed.⁷ The apparatus consisted simply of a large, degassed 5 l. fused quartz bulb, modified with several attached, heated "reactor" zones of varying dimensions (cylindrically symmetric) distributed over the surface. The bulb was connected to a gas handling system and could

also be heated to alter the input distribution. Entrance area to a reactor was restricted to $\gtrsim 1/100$ of the bulb area. Dimensions of the reactors used so far provided mean wall collision numbers, \bar{m} , of 2, 4.5, 10.5 and 22.3, per encounter. For a given \bar{m} , the distribution of wall collisions was found by Monte Carlo calculation.^{7,8} Molecules that returned to the bulb were cooled by collisions before another encounter with the reactor. Any value of \bar{m} from 1 to 10^4 is readily attained; the technique is termed the Variable Encounter Method.

We have applied the method to the structural isomerization of 1,1-cyclo-propane-d₂ activated by collisions at a seasoned surface_A. This system is notably free from complications. Brief pre-exposure of the oxidized silica wall to substrate provided adequate seasoning and reproducible results. To perform a run, one reactor was heated to a desired temperature, the bulb (at 375 K) was pumped to a stable residual pressure of $< 10^{-6}$ torr, and substrate was admitted at a pressure of $1-2 \times 10^{-4}$ torr and left for a suitable time. A flow mode was also possible. Analysis was by gas chromatography; percent reaction varied from 1 - 50%.

Figure 1 shows experimental temperature plots of the average reaction probability per collision, \bar{P}_c , for the four reactors. The data were compared to stochastic calculations using two forms of P , gaussian and exponential. Several models were computed in which the average downjump step size, $\langle \Delta E \rangle$, varied as E^x , $x = 0, 1/2, 1$. The gaussian form (with $x = 0, 1/2$) tentatively give the best (cf. refs. 6 and 9). fits_A. Some numerical values of $\langle \Delta E \rangle$ for the case $x = 0$, are given in the figure (use of $x = 1/2, 1$ leads, of course, to larger values of $\langle \Delta E \rangle$ at the level of energy E_0). Once P has been selected, $\bar{\tau}$ follows immediately.^{3,4}

Figure 2a illustrates the variation of \bar{P}_c with \bar{m} at constant temperature. Figure 2b shows the sequential variation of P_c vs collision number, calculated

from the flat gaussian model.

We find: a) For the temperatures covered, accommodation to a steady state with the wall takes place in a small number of collisions $\sim 8-15$ (Fig. 2b), the number increasing with temperature; this accounts for the findings made in low pressure pyrolysis studies¹⁰ that \bar{m} values $\gtrsim 80-100$ provide temperature accommodation; at lower temperatures, especially, wall efficiency approaches strong collider behavior and $\bar{\tau}$ is quasi-independent of the input distribution (bulb temperature). b) Wall collisions are more efficient than gas phase collisions; the surface seasoned with a film of "polycyclopropane" is characterized by down step sizes $\langle \Delta E \rangle$ of 4,000 and $3,400 \text{ cm}^{-1}$, ($x = 0$), at 775 K and 975 K, respectively; these compare with the lesser values, $\sim 3,500$ and $1,900 \text{ cm}^{-1}$, measured previously¹² for binary gas phase cyclopropane-d₂ collisions; c) The present data thus support the previous finding that energy transfer collision efficiency declines at higher temperatures for this polyatomic unimolecular system. Dove and co-workers¹³ have estimated an "incubation" period of $\sim 3,000$ collisions for the shock-heated triatomic N₂O system above 2000K.

Features such as surface structure, substrate structure and level of E_0 are subjects for future investigation by this technique.

References

- a) This work was supported by the Office of Naval Research and in part by the National Science Foundation.
 - b) UNIDO Fellow. University of Szeged, Szeged, Hungary.
1. R. J. Rubin and K. E. Shuler, *J. Chem. Phys.* 25, 59,68 (1956).
 2. E. W. Montroll and K. E. Shuler, *Adv. Chem. Phys.* 1, 361 (1958).
 3. S. K. Kim, *J. Chem. Phys.* 28, 1057 (1958).
 4. B. Widom, *J. Chem. Phys.* 31, 1387 (1959); 34, 2050 (1961).
 5. Y. N. Lin and B. S. Rabinovitch, *J. Phys. Chem.* 74, 3151 (1970).
 6. S. C. Chan, B. S. Rabinovitch, J. T. Bryant, L. D. Spicer, T. Fujimoto, Y. N. Lin and S. P. Pavlou, *J. Phys. Chem.* 74, 3160 (1970); D. C. Tardy and B. S. Rabinovitch, *Chem. Revs.* 77, 369 (1977).
 7. B. D. Barton, Ph.D. Thesis, University of Washington, in preparation.
 8. D. H. Davis, *J. Appl. Phys.* 31, 1169 (1960).
 9. G. Kohlmaier and B. S. Rabinovitch, *J. Chem. Phys.*, 38, 1692, 1709 (1963).
 10. G. N. Spokes and S. W. Benson, *J. Am. Chem. Soc.*, 89, 2527 (1967); K. D. King, D. M. Golden, G. N. Spokes and W. S. Benson, *Int. J. Chem. Kin.* 3, 411 (1971).
 11. Unpublished work on cyclopropane (ref. 7) even indicates somewhat higher values.
 12. I. E. Klein, B. S. Rabinovitch, and K. H. Jung, *J. Chem. Phys.* 67, 3833 (1977). I. E. Klein and B. S. Rabinovitch, *J. Phys. Chem.*, 82, 243 (1978).
 13. J. E. Dove, W. S. Nip and H. Teitelbaum, *Intern. Sympos. Combust.* 15, 903 (1974).

Figure Captions

1. Plots of experimental values of \bar{P}_c vs T for each of the four reactors. Solid curves are calculated fits for a gaussian model ($x = 0$) of the energy transfer probabilities for each value of m , and include the limiting case, $m = \infty$. The best-fit values of $\langle \Delta E \rangle$ (cm^{-1}) are given at several temperatures.
2. A) Illustrative plots of experimental values of \bar{P}_c vs m at 823K, 973K and 1123K. Solid curves are calculated for respective values of $\langle \Delta E \rangle$ of 3800, 3400 and 3050 cm^{-1} . The ordinate scales are arbitrary and are normalized to $\bar{P}_c = 1$ for $m = \infty$, at each temperature.
B) Illustrative histograms of the calculated sequential probability $P_c(n)$ vs n , the number of consecutive collisions, for the temperatures and values of $\langle \Delta E \rangle$ used in A).




Fig 1 Kelley et al




Fig 2. Kelley et al